



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 7 :</b> <b>C07K 14/435, C12N 1/00, 1/15, 1/21, 5/10, 15/12, 15/63</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/34318</b> <b>(43) International Publication Date:</b> 15 June 2000 (15.06.00)
<b>(21) International Application Number:</b> PCT/US99/29291 <b>(22) International Filing Date:</b> 10 December 1999 (10.12.99)  <b>(30) Priority Data:</b> 09/210,330 11 December 1998 (11.12.98) US Not furnished 9 December 1999 (09.12.99) US  <b>(71) Applicant:</b> CLONTECH LABORATORIES, INC. [US/US]; 1020 East Meadow Drive, Palo Alto, CA 94303 (US).  <b>(72) Inventors:</b> LUKYANOY, Sergey, Anatolievich; ul. Golubinskaya 13/1-161, Moscow (RU). FRADKOV, Arcady, Fedorovich; ul. Dnepropetrovskaya, 35/2-14, Moscow, 113570 (RU). LABAS, Yulii Aleksandrovich; ul. Generala Tyuleneva, 35-416, Moscow, 117465 (RU). MATZ, Mikhail Vladimirovich; ul. Teplii stan, 7/2-28, Moscow, 117465 (RU). JIANG, Xin; 1133 Rajkovich Way, San Jose, CA 95120 (DE). DUONG, Tommy; 2491 Glen Elm Way, San Jose, CA 95148 (US).  <b>(74) Agent:</b> ADLER, Benjamin, A.; McGregor & Adler, 8011 Candle Ln., Houston, TX 77071 (US).		<b>(81) Designated States:</b> JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> FLUORESCENT PROTEINS FROM NON-BIOLUMINESCENT SPECIES OF CLASS ANTHOZOA, GENES ENCODING SUCH PROTEINS AND USES THEREOF  <b>(57) Abstract</b>  The present invention is directed to novel fluorescent proteins from non-bioluminescent organisms from the Class Anthozoa. Also disclosed are cDNAs encoding the fluorescent proteins.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**FLUORESCENT PROTEINS FROM NON-BIOLUMINESCENT SPECIES  
OF CLASS ANTHOZOA, GENES ENCODING SUCH PROTEINS AND  
USES THEREOF**

10

**BACKGROUND OF THE INVENTION**

Cross-reference to Related Application

This is a divisional application of U.S.S.N. 09/210,330 filed  
15 on December 11, 1998.

Field of the Invention

This invention relates to the field of molecular biology.  
More specifically, this invention relates to novel fluorescent proteins,  
20 cDNAs encoding the proteins and uses thereof.

Description of the Related Art

Fluorescence labeling is a particularly useful tool for  
marking a protein, cell, or organism of interest. Traditionally, a  
25 protein of interest is purified, then covalently conjugated to a  
fluorophore derivative. For *in vivo* studies, the protein-dye complex is  
then inserted into cells of interest using micropipetting or a method of  
reversible permeabilization. The dye attachment and insertion steps,  
however, make the process laborious and difficult to control. An

alternative method of labeling proteins of interest is to concatenate or fuse the gene expressing the protein of interest to a gene expressing a marker, then express the fusion product. Typical markers for this method of protein labeling include  $\beta$ -galactosidase, firefly luciferase and bacterial luciferase. These markers, however, require exogenous substrates or cofactors and are therefore of limited use for *in vivo* studies.

A marker that does not require an exogenous cofactor or substrate is the green fluorescent protein (GFP) of the jellyfish *Aequorea victoria*, a protein with an excitation maximum at 395 nm, a second excitation peak at 475 nm and an emission maximum at 510 nm. GFP is a 238-amino acid protein, with amino acids 65-67 involved in the formation of the chromophore.

Uses of GFP for the study of gene expression and protein localization are discussed in detail by Chalfie et al. in *Science* 263 (1994), 802-805, and Heim et al. in *Proc. Nat. Acad. Sci.* 91 (1994), 12501-12504. Additionally, Rizzuto et al. in *Curr. Biology* 5 (1995), 635-642, discuss the use of wild-type GFP as a tool for visualizing subcellular organelles in cells, while Kaether and Gerdes in *Febs Letters* 369 (1995), 267-271, report the visualization of protein transport along the secretory pathway using wild-type GFP. The expression of GFP in plant cells is discussed by Hu and Cheng in *Febs Letters* 369 (1995), 331-334, while GFP expression in *Drosophila* embryos is described by Davis et al. in *Dev. Biology* 170 (1995), 726-729.

Crystallographic structures of wild-type GFP and the mutant GFP S65T reveal that the GFP tertiary structure resembles a barrel (Ormö et al., *Science* 273 (1996), 1392-1395; Yang, et al., *Nature Biotechnol* 14 (1996), 1246-1251). The barrel consists of beta sheets in a compact structure, where, in the center, an alpha helix containing

the chromophore is shielded by the barrel. The compact structure makes GFP very stable under diverse and/or harsh conditions such as protease treatment, making GFP an extremely useful reporter in general. However, the stability of GFP makes it sub-optimal for  
5 determining short-term or repetitive events.

A great deal of research is being performed to improve the properties of GFP and to produce GFP reagents useful and optimized for a variety of research purposes. New versions of GFP have been developed, such as a "humanized" GFP DNA, the protein product of  
10 which has increased synthesis in mammalian cells (Haas, et al., *Current Biology* 6 (1996), 315-324; Yang, et al., *Nucleic Acids Research* 24 (1996), 4592-4593). One such humanized protein is "enhanced green fluorescent protein" (EGFP). Other mutations to GFP have resulted in blue-, cyan- and yellow-green light emitting versions. Despite the great  
15 utility of GFP, however, other fluorescent proteins with properties similar to or different from GFP would be useful in the art. Novel fluorescent proteins result in possible new colors, or produce pH-dependent fluorescence. Other benefits of novel fluorescent proteins include fluorescence resonance energy transfer (FRET) possibilities  
20 based on new spectra and better suitability for larger excitation.

The prior art is deficient in novel fluorescent proteins wherein the DNA coding sequences are known. The present invention fulfills this long-standing need in the art.

25

### SUMMARY OF THE INVENTION

The present invention is directed to DNA sequences encoding fluorescent proteins selected from the group consisting of:

(a) an isolated DNA from an organism from the Class Anthozoa which encodes a fluorescent protein; (b) an isolated DNA which hybridizes to the isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code and that encodes a fluorescent protein. Preferably, the DNA is isolated from a non -bioluminescent organism from Class Anthozoa. More preferably, the DNA has the sequence selected from the group consisting of SEQ ID Nos. 55, 57-60, and the fluorescent protein has the amino acid sequence shown in SEQ ID No. 56.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention in a recombinant cell comprising said DNA and regulatory elements necessary for expression of the DNA in the cell. Preferably, the DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56.

In still another embodiment of the present invention, there is provided a host cell transfected with a vector of the present invention, such that the host cell expresses a fluorescent protein. Preferably, the cell is selected from the group consisting of bacterial cells, mammalian cells, plant cells and insect cells.

The present invention is also directed to an isolated and purified fluorescent protein coded for by DNA selected from the group consisting of: (a) isolated DNA from an organism from Class Anthozoa which encodes a fluorescent protein; (b) isolated DNA which hybridizes to the isolated DNA of (a) and which encodes a fluorescent protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code, and which

encodes a fluorescent protein. Preferably, the protein has the amino acid sequence shown in SEQ ID No. 56.

The present invention is also directed to a DNA sequence encoding a fluorescent protein selected from the group consisting of:

5 (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated DNA which hybridizes to isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of

10 (a) and (b) in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein. Preferably, the organism is from Sub-class Zoantharia, Order Zoanthidea. More preferably, the organism is from Sub-order Brachynermida. Even more preferably, the organism is from Family Zoanthidae, Genus Zoanthus. Most

15 particularly, the present invention is drawn to a novel fluorescent protein from *Zoanthus* sp., zFP506.

The present invention is further directed to an amino acid sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by

20 means of hybridization, wherein the amino acid sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14. Preferably, such an oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16.

Other and further aspects, features, and advantages of the

25 present invention will be apparent from the following description of the presently preferred embodiments of the invention given for the purpose of disclosure.

## BRIEF DESCRIPTION OF THE DRAWINGS

**Figure 1** shows the modified strategy of 3'-RACE used to isolate the target fragments. Sequences of the oligonucleotides used are shown in Table 2. Dp1 and Dp2 are the degenerate primers used in the first and second PCR, respectively (see Tables 3 and 4 for the sequences of degenerate primers). In the case of *Zoanthus sp.*, the first degenerate primer used was NGH (SEQ ID No. 4), and the second degenerate primer used was GEGa (SEQ ID No. 6).

**Figure 2** shows the excitation and emission spectrum of the novel fluorescent protein from *Zoanthus sp.*, zFP506.

**Figure 3** shows functional analysis of destabilized zFP506. It demonstrates that fusion of MODC domain to the zFP506 slightly decreases the fluorescent intensity compared to zFP506 itself and that transient transfection of destabilized zFP506 decreases the fluorescent intensity by 50% after 4-hour treatment with protein synthesis inhibitor cycloheximide (**Figure 3B**) vs. control (**Figure 3A**).

## DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "GFP" refers to the basic green fluorescent protein from *Aequorea victoria*, including prior art versions of GFP engineered to provide greater fluorescence or fluoresce in different colors. The sequence of *Aequorea victoria* GFP (SEQ ID No. 54) has been disclosed in Prasher et al., *Gene* 111 (1992), 229-33.

As used herein, the term "EGFP" refers to mutant variant of GFP having two amino acid substitutions: F64L and S65T (Heim et al., *Nature* 373 (1995), 663-664). The term "humanized" refers to changes



made to the GFP nucleic acid sequence to optimize the codons for expression of the protein in human cells (Yang et al., *Nucleic Acids Research* 24 (1996), 4592-4593).

As used herein, the term "NFP" refers to novel fluorescent protein, and the term "GNFP" refers to green novel fluorescent protein. Specifically, "GNFP" refers to zFP506.

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Maniatis, Fritsch & Sambrook, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins eds. (1985)); "Transcription and Translation" (B.D. Hames & S.J. Higgins eds. (1984)); "Animal Cell Culture" (R.I. Freshney, ed. (1986)); "Immobilized Cells and Enzymes" (IRL Press, (1986)); B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in either single stranded form or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes.

A DNA "coding sequence" is a DNA sequence which is transcribed and translated into a polypeptide in vivo when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and synthetic DNA sequences. A polyadenylation signal and transcription termination sequence may be located 3' to the coding sequence.

As used herein, the term "hybridization" refers to the process of association of two nucleic acid strands to form an antiparallel duplex stabilized by means of hydrogen bonding between residues of the opposite nucleic acid strands.

The term "oligonucleotide" refers to a short (under 100 bases in length) nucleic acid molecule.

"DNA regulatory sequences", as used herein, are transcriptional and translational control sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for and/or regulate expression of a coding sequence in a host cell.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a

transcription initiation site, as well as protein binding domains responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes. Various promoters, including inducible promoters, may be used to  
5 drive the various vectors of the present invention.

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" or "transfected" by  
10 exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to  
15 eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter  
20 cells containing the transforming DNA. A "clone" is a population of cells derived from a single cell or common ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

A "heterologous" region of the DNA construct is an  
25 identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. In another example,

heterologous DNA includes coding sequence in a construct where portions of genes from two different sources have been brought together so as to produce a fusion protein product. Allelic variations or naturally-occurring mutational events do not give rise to a  
5 heterologous region of DNA as defined herein.

As used herein, the term "reporter gene" refers to a coding sequence attached to heterologous promoter or enhancer elements and whose product may be assayed easily and quantifiably when the construct is introduced into tissues or cells.

10 The amino acids described herein are preferred to be in the "L" isomeric form. The amino acid sequences are given in one-letter code (A: alanine; C: cysteine; D: aspartic acid; E: glutamic acid; F: phenylalanine; G: glycine; H: histidine; I: isoleucine; K: lysine; L: leucine; M: methionine; N: asparagine; P: proline; Q: glutamine; R: arginine; S:  
15 serine; T: threonine; V: valine; W: tryptophane; Y: tyrosine; X: any residue).  $\text{NH}_2$  refers to the free amino group present at the amino terminus of a polypeptide.  $\text{COOH}$  refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J Biol. Chem.*, 243 (1969), 3552-  
20 59 is used.

The present invention is directed to an isolated DNA selected from the group consisting of: (a) isolated DNA from an organism from the Class Anthozoa which encodes a fluorescent protein; (b) isolated DNA which hybridizes to isolated DNA of (a) and  
25 which encodes a fluorescent protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code, and which encodes a fluorescent protein. Preferably, the DNA has the sequence selected from the group consisting of SEQ ID Nos. 55, 57-60, and the fluorescent protein has the

amino acid sequence shown in SEQ ID No. 56. More preferably, the DNA is non-humanized or humanized zFP506 or N65M.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention in a recombinant cell comprising said DNA and regulatory elements necessary for expression of the DNA in the cell. Specifically, the DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56. Preferably, the vector is constructed by amplifying the DNA and then inserting the amplified DNA into EGFP-N1 backbone, or by fusing different mouse ODC degradation domains such as d1, d2 and d376 to the C-terminal of the DNA and then inserting the fusion DNA into EGFP-N1 backbone.

In still another embodiment of the present invention, there is provided a host cell transfected with the vector of the present invention, which expresses a fluorescent protein of the present invention. Preferably, the cell is selected from the group consisting of bacterial cells, mammalian cells, plant cells, insect cells and yeast cells. A representative example of mammalian cell is HEK 293 cell and an example of bacterial cell is an *E. coli* cell.

The present invention is also directed to a DNA sequence encoding a fluorescent protein selected from the group consisting of: (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated DNA which hybridizes to isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein. Preferably, the organism is from Sub-class Zoantharia, Order Zoanthidea. More preferably, the

organism is from Sub-order Brachycnemina. Even more preferably, the organism is from Family Zoanthidae, Genus Zoanthus.

The present invention is also directed to an isolated and purified fluorescent protein coded for by DNA selected from the group consisting of: (a) an isolated protein encoded by a DNA which encodes a fluorescent protein wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated protein encoded by a DNA which hybridizes to isolated DNA of (a); and (c) an isolated protein encoded by a DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code. Preferably, the isolated and purified fluorescent protein is zFP506.

The present invention is further directed to an amino acid sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by means of hybridization, wherein the amino acid sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14. Preferably, such an oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16 and is used as a primer in polymerase chain reaction. Alternatively, it can be used as a probe for hybridization screening of the cloned genomic or cDNA library.

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion.

**EXAMPLE 1****Biological Material**

Novel fluorescent proteins were identified from several  
5 genera of Anthozoa which do not exhibit any bioluminescence but have  
fluorescent color as observed under usual white light or ultraviolet  
light. Six species were chosen (see Table 1).

**TABLE 1**Anthozoa Species Used in This Study

Species	Area of Origination	Fluorescent Color
Anemonia majano	Western Pacific	bright green tentacle tips
Clavularia sp.	Western Pacific	bright green tentacles and oral disk
Zoanthus sp.	Western Pacific	green-yellow tentacles and oral disk
Discosoma sp. "red"	Western Pacific	orange-red spots oral disk
Discosoma striata	Western Pacific	blue-green stripes on oral disk
Discosoma sp. "magenta"	Western Pacific	faintly purple oral disk
Discosoma sp. "green"	Western Pacific	green spots on oral disk
Anemonia sulcata	Mediterranean	purple tentacle tips



**EXAMPLE 2****cDNA Preparation**

Total RNA was isolated from the species of interest according to the protocol of Chomczynski and Sacchi (Chomczynski P., et al., *Anal. Biochem.* 162 (1987), 156-159). First-strand cDNA was synthesized starting with 1-3 µg of total RNA using SMART PCR cDNA synthesis kit (CLONTECH) according to the provided protocol with the only alteration being that the "cDNA synthesis primer" provided in the kit was replaced by the primer TN3 (5'-CGCAGTCGACCG(T)<sub>13</sub>, SEQ ID No. 1) (Table 2). Amplified cDNA samples were then prepared as described in the protocol provided except the two primers used for PCR were the TS primer (5'-AAGCAGTGGTATCAACGCAGAGT, SEQ ID No. 2) (Table 2) and the TN3 primer (Table 2), both in 0.1 µM concentration. Twenty to twenty-five PCR cycles were performed to amplify a cDNA sample. The amplified cDNA was diluted 20-fold in water and 1 µl of this dilution was used in subsequent procedures.

### Oligos Used in cDNA Synthesis and RACE

25

**EXAMPLE 3**Oligo Design

To isolate fragments of novel fluorescent protein cDNAs,  
5 PCR using degenerate primers was performed. Degenerate primers  
were designed to match the sequence of the mRNAs in regions that  
were predicted to be the most invariant in the family of fluorescent  
proteins. Four such stretches were chosen (Table 3) and variants of  
degenerate primers were designed. All such primers were directed to  
10 the 3'-end of mRNA. All oligos were gel-purified before use. Table 2  
shows the oligos used in cDNA synthesis and RACE.

**TABLE 3**

Key Amino Acid Stretches and Corresponding Degenerate Primers Used  
for Isolation of Fluorescent Proteins

5

Stretch Position according to A. victoria GFP (7)	Amino Acid Sequence of the Key Stretch	Degenerated Primer Name and Sequence
20-25	GXVNGH (SEQ ID No. 3)	NGH: 5'- GA(C,T) GGC TGC GT(A,T,G,C) AA(T,C) GG(A,T,G) CA (SEQ ID No. 4)
31-35	GEGEG (SEQ ID No. 5)  GEGNG (SEQ ID No. 8)	GEGa: 5'- GTT ACA GGT GA(A,G) GG(A,C) GA(A,G) GG (SEQ ID No. 6) GEGb: 5'- GTT ACA GGT GA(A,G) GG(T,G) GA(A,G) GG (SEQ ID No. 7) GNGa: 5'- GTT ACA GGT GA(A,G) GG(A,C) AA(C,T) GG (SEQ ID No. 9) GNGb: 5'- GTT ACA GGT GA(A,G) GG(T,G) AA(C,T) GG (SEQ ID No. 10)
127-131	GMNFP (SEQ ID No. 11) GVNFP (SEQ ID No. 12)	NFP: 5' TTC CA(C,T) GGT (G,A)TG AA(C,T) TT(C,T) CC (SEQ ID NO. 13)
134-137	GPVM (SEQ ID No. 14)	PVMa: 5' CCT GCC (G,A)A(C,T) GGT CC(A,T,G,C) GT(A,C) ATG (SEQ ID NO. 15) PVMb: 5' CCT GCC (G,A)A(C,T) GGT CC(A,T,G,C) GT(G,T) ATG (SEQ ID NO. 16)

**EXAMPLE 4**Isolation of 3'-cDNA Fragments of nEPs

The modified strategy of 3'-RACE was used to isolate the target fragments (see Figure 1). The RACE strategy involved two consecutive PCR steps. The first PCR step involved a first degenerate primer (Table 4) and the T7-TN3 primer (SEQ ID No. 17) which has a 3' portion identical to the TN3 primer used for cDNA synthesis (for sequence of T7-TN3, Table 2). The reason for substituting the longer T7-TN3 primer in this PCR step was that background amplification which occurred when using the shorter TN3 primer was suppressed effectively, particularly when the T7-TN3 primer was used at a low concentration (0.1  $\mu$ M) (Frohman et al., (1998) *PNAS USA*, 85, 8998-9002). The second PCR step involved the TN3 primer (SEQ ID No. 1, Table 2) and a second degenerate primer (Table 4).

**TABLE 4**

Combinations of Degenerate Primers for First and Second PCR Resulting in Specific Amplification of 3'-Fragments of nFP cDNA

Species	First Degenerate Primer	Second Degenerate Primer
Anemonia majano	NGH (SEQ ID No. 4)	GNGb – (SEQ ID No. 10)
Clavularia sp.	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6)
Zoanthus sp.	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6)
Discosoma sp. "red"	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6), NFP (SEQ ID No. 13) or PVMb (SEQ ID No. 16)
Discosoma striata	NGH (SEQ ID No. 4)	NFP (SEQ ID No. 13)
Anemonia sulcata	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6) or NFP (SEQ ID No. 13)

5

The first PCR reaction was performed as follows: 1 µl of 20-fold  
 10 dilution of the amplified cDNA sample was added into the reaction  
 mixture containing 1X Advantage KlenTaq Polymerase Mix with  
 provided buffer (CLONTECH), 200 µM dNTPs, 0.3 µM of first degenerate

primer (Table 4) and 0.1  $\mu$ M of T7-TN3 (SEQ ID No. 17) primer in a total volume of 20  $\mu$ l. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 1 cycle for 95°C, 10 sec.; 55°C, 1 min.; 72°C, 40 sec; 24 cycles for 95°C, 10 sec.; 62°C, 30 sec.; 72°C, 40 sec. The reaction was then diluted 20-fold in water and 1  $\mu$ l of this dilution was added to a second PCR reaction, which contained 1X Advantage KlenTaq Polymerase Mix with the buffer provided by the manufacturer (CLONTECH), 200  $\mu$ M dNTPs, 0.3  $\mu$ M of the second degenerate primer (Table 4) and 0.1  $\mu$ M of TN3 primer. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 1 cycle for 95°C, 10 sec.; 55°C (for GEG/GNG or PVM) or 52°C (for NFP), 1 min.; 72°C, 40 sec; 13 cycles for 95°C, 10sec.; 62°C (for GEG/GNG or PVM) or 58°C (for NFP), 30 sec.; 72°C, 40 sec. The product of PCR was cloned into PCR-Script vector (Stratagene) according to the manufacturer's protocol.

Different combinations of degenerate primers were tried in the first and second PCR reactions on the DNA from each species until a combination of primers was found that resulted in specific amplification--meaning that a pronounced band of expected size (about 650-800 bp for NGH and GEG/GNG and 350-500 bp for NFP and PVM--sometimes accompanied by a few minor bands) was detected on agarose gel after two PCR reactions. The primer combinations of choice for different species of the Class Anthozoa are listed in Table 4. Some other primer combinations also resulted in amplification of fragments of correct size, but the sequence of these fragments showed no homology to the other fluorescent proteins identified or to *Aequorea victoria* GFP.

**EXAMPLE 5**Obtaining Full-Length cDNA Copies

Upon sequencing the obtained 3'-fragments of novel  
5 fluorescent protein cDNAs, two nested 5'-directed primers were  
synthesized for cDNA (Table 5), and the 5' ends of the cDNAs were  
then amplified using two consecutive PCRs. In the next PCR reaction,  
the novel approach of "step-out PCR" was used to suppress background  
amplification. The step-out reaction mixture contained 1x Advantage  
10 KlenTaq Polymerase Mix using buffer provided by the manufacturer  
(CLONTECH), 200  $\mu$ M dNTPs, 0.2  $\mu$ M of the first gene-specific primer  
(see Table 5), 0.02  $\mu$ M of the T7-TS primer (SEQ ID No. 18), 0.1  $\mu$ M of  
T7 primer (SEQ ID No. 19) and 1  $\mu$ l of the 20-fold dilution of the  
amplified cDNA sample in a total volume of 20  $\mu$ l. The cycling profile  
15 was (Hybaid OmniGene Thermocycler, tube control mode): 23-27  
cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of  
amplification was diluted 50-fold in water and one  $\mu$ l of this dilution  
was added to the second (nested) PCR. The reaction contained 1X  
Advantage KlenTaq Polymerase Mix with provided buffer (CLONTECH),  
20 200  $\mu$ M dNTPs, 0.2  $\mu$ M of the second gene-specific primer and 0.1  $\mu$ M  
of TS primer (SEQ ID No. 2) in a total volume of 20  $\mu$ l. The cycling  
profile was (Hybaid OmniGene Thermocycler, tube control mode): 12  
cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of  
amplification was then cloned into pAtlas vector (CLONTECH) according  
25 to the manufacturer's protocol.



**TABLE 5**
Gene-Specific Primers Used for 5'-RACE

Species	First Primer	Second (Nested) Primer
Anemonia majano	5'-GAAATAGTCAGGCATACTGGT (SEQ ID No. 20)	5'-GTCAGGCATAC TGGTAGGAT (SEQ ID No. 21)
Clavularia sp.	5'-CTTGAAATAGTCTGCTATATC (SEQ ID No. 22)	5'-TCTGCTATATC GTCTGGGT (SEQ ID No. 23)
Zoanthus sp.	5'- GTTCTTGAAATAGTCTACTATGT (SEQ ID No. 24)	5'-GTCTACTATGTCTT GAGGAT (SEQ ID No. 25)
Discosoma sp. "red"	5'-CAAGCAAATGGCAAAGGTC (SEQ ID No. 26)	5'-CGGTATTGTGGCC TTCGTA (SEQ ID No. 27)
Discosoma striata	5'-TTGTCTTCTTCTGCACAAC (SEQ ID No. 28)	5'-CTGCACAACGG GTCCAT (SEQ ID No. 29)
Anemonia sulcata	5'-CCTCTATCTTCATTTCCTGC (SEQ ID No. 30)	5'-TATCTTCATTTCCT GCGTAC (SEQ ID No. 31)
Discosoma sp. "magenta"	5'-TTCAGCACCCCATCACGAG (SEQ ID No. 32)	5'-ACGCTCAGAGCTG GGTTCC (SEQ ID No. 33)
Discosoma sp. "green"	5'-CCCTCAGCAATCCATCACGTTC (SEQ ID No. 34)	5'-ATTATCTCAGTGGA TGGTTC (SEQ ID No. 35)

**EXAMPLE 6**Expression of nFP in *E. coli*

5 To prepare a DNA construct for novel fluorescent protein expression, two primers were synthesized for each cDNA: a 5'-directed "downstream" primer with the annealing site located in the 3'-UTR of the cDNA and a 3'-directed "upstream" primer corresponding to the site of translation start site (not including the first ATG codon) (Table 10 6). Primers with SEQ ID Nos. 41 and 42 were the primers used to prepare the zFP506 DNA. Both primers had 5'-heels coding for a site for a restriction endonuclease; in addition, the upstream primer was designed so as to allow the cloning of the PCR product into the pQE30 vector (Qiagen) in such a way that resulted in the fusion of reading 15 frames of the vector-encoded 6xHis-tag and nFP. The PCR was performed as follows: 1 µl of the 20-fold dilution of the amplified cDNA sample was added to a mixture containing 1x Advantage KlenTaq Polymerase Mix with buffer provided by the manufacturer (CLONTECH), 200 µM dNTPs, 0.2 µM of upstream primer and 0.2 µM of downstream 20 primer, in a final total volume of 20 µl. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 23-27 cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of this amplification step was purified by phenol-chlorophorm extraction and ethanol precipitation and then cloned into pQE30 vector using restriction 25 endonucleases corresponding to the primers' sequence according to standard protocols.

All plasmids were amplified in XL-1 blue *E. coli* and purified by plasmid DNA miniprep kits (CLONTECH). The recombinant clones were selected by colony color, and grown in 3 ml of LB medium.

(supplemented with 100  $\mu\text{g/ml}$  of ampicillin) at 37°C overnight. 100  $\mu\text{l}$  of the overnight culture was transferred into 200 ml of fresh LB medium containing 100  $\mu\text{g/ml}$  of ampicillin and grown at 37°C, 200 rpm up to OD<sub>600</sub> 0.6-0.7. 1 mM IPTG was then added to the culture and 5 incubation was allowed to proceed at 37°C for another 16 hours. The cells were harvested and recombinant protein, which incorporated 6x His tags on the N-terminus, was purified using TALON™ metal-affinity resin according to the manufacturer's protocol (CLONTECH).

**TABLE 6**

Primers Used to Obtain Full Coding Region of nFPs for Cloning into Expression Construct

Species	Upstream Primer	Downstream Primer
Anemonia majano	5' -acatggatccgctctttcaaca agttatc (SEQ ID No. 36) BamHI	5'-tagtactcgcgagcttattcgta tttcagtgaatc (SEQ ID No. 37) XhoI
Clavularia sp.	L: 5'-acatggatccaacatttttga gaaacg (SEQ ID No. 38) BamHI S: 5'-acatggatccaaagctctaacc accatg (SEQ ID No. 39) BamHI	5'-tagtactcgcgagcaacacaa accctcagacaa (SEQ ID No. 40) XhoI
Zoanthus sp.	5'- acatggatccgctcagtcacaaag cacggt (SEQ ID No. 41) BamHI	5'-tagtactcgcgaggttggaactacat tcttatca (SEQ ID No. 42) XhoI
Discosoma sp. "red"	5'- acatggatccaggtcttccaagaat gttatc (SEQ ID No. 43) BamHI	5'-tagtactcgcgaggagccaagttc agcctta (SEQ ID No. 44) XhoI
Discosoma striata	5'- acatggatccagttggtccaagagtgtg (SEQ ID No. 45) BamHI	5'-tagcgcgagctctatcatgcctc gtcacct (SEQ ID No. 46) SacI
Anemonia sulcata	5'- acatggatccgcttccttttaagaagact (SEQ ID No. 47) BamHI	5'-tagtactcgcgagtccttgggagc ggcttg (SEQ ID No. 48) XhoI
Discosoma sp. "magenta"	5'- acatggatccagttgttccaagaatgtgat (SEQ ID No. 49) BamHI	5'-tagtactcgcgaggccattacg ctaac (SEQ ID No. 50) XhoI
Discosoma sp. "green"	5'-acatggatccagtgcaacttaagaagaaatg (SEQ ID No. 51)	5'-tagtactcgcgagattcggtttaat gccttg (SEQ ID No. 52)

5

**EXAMPLE 7**Novel Fluorescent Proteins and cDNAs Encoding the Proteins

One of the full-length cDNAs encoding fluorescent proteins  
 5 found is described herein (zFP506). The nucleic acid sequence and  
 deduced amino acid sequence are SEQ ID Nos. 55 and 56, respectively.  
 The spectral properties of zFP506 is listed in Table 7, and the emission  
 and excitation spectra for the ZFP506 is shown in Figure 2.

10

**TABLE 7**Spectral Properties of the Isolated zFP506 nFP

	Species:	Zoanthus sp.	Max. Extinction Coefficient:	35,600
15	nFP Name:	zFP506	Quantum Yield	0.63
	Absorbance Max. (nm):	496	Relative Brightness:*	1.02
20	Emission Max. (nm):	506		

\*relative brightness is extinction coefficient multiplied by quantum  
 yield divided by the same value for *A. victoria* GFP.

25

**EXAMPLE 8**Construction of zFP506 Mutant

One mutant of zFP506 was generated, N65M. Compared  
 30 with wild type zFP506, N65M has the mutation of from "AAC" to "ATG"  
 which results in the corresponding amino acid change from Asparagine

(N) to Methionine (M) at the position of 65. The spectral properties of N65M are listed in Table 8.

5

**TABLE 8**Spectral Properties of the Isolated N65M

10	Species:	Zoanthus sp.	Max. Extinction Coefficient:	62,000
	nFP Name:	N65M	Quantum Yield	0.63
15	Absorbance Max. (nm):	496	Relative Brightness:*	1.78
	Emission Max. (nm):	506		

\*relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

20

**EXAMPLE 9**Construction and Functional Analysis of Vectors

Non-humanized zFP506 DNA was amplified via PCR and  
25 reconstructed into EGFP-N1 backbone. This vector has the same  
multiple cloning sites as EGFP-N1.

Functional test of the generated vector was performed by  
transient transfection in 293 cells. 24 hours post transfection,  
expression of zFP506 was examined under fluorescent microscope.  
30 zFP506 showed good fluorescent intensity and comparable to EGFP-N1.

**EXAMPLE 10**Generation of Destabilized zFP506 Vectors as Transcription Reporters

5                Since zFP506 is very stable, it is necessary to generate destabilized versions of zFP506 in order to observe the rapid turnover of the protein. By using the same technology for destabilized EGFP, two destabilized zFP506 vectors were constructed by fusing mouse ODC degradation domain to the C-terminal of zFP506. The d1 version of  
10 destabilized zFP506 (SEQ ID No. 57) has three E to A mutations within MODC degradation domain comparing to d2 version (SEQ ID No. 58), therefore result in a shorter half-life of the protein to which MODC degradation domain fused to. Destablized d1zFP506 and d2zFP506 were constructed in EGFP-N1 backbone

15

**EXAMPLE 10**Functional Analysis of Destabilized zFP506

20                Wildtype d1zFP506 was transiently transfected into 293 cells. 24 hours after transfection, CHX was added to stop protein synthesis. After 4 hour treatment, cells were examined under fluorescent microscope. It shows that fusion of MODC domain to the zFP506 slightly decreases the fluorescent intensity compared to zFP506  
25 itself. After 4 hour treatment, there is 50% fluorescent intensity decrease (Figure 3B vs. Figure 3A).

**EXAMPLE 11****Application of Destabilized dlzFP506 as Transcription Reporters**

Destabilized dlzFP506 was constructed into pCRE-dlGNFP  
5 and pNF- $\kappa$ B-dlGNFP vectors. Its expression was regulated under cAMP  
response element (CRE) or NF- $\kappa$ B response element, respectively. These  
vectors were transiently transfected into 293 cells, and 24 hours post  
transfection, the expression of dlGNFP was induced by Forskolin or  
TNF- $\alpha$ . 6 hours after induction, the culture was analysed by FACS. CRE-  
10 dlGNFP showed 7 fold of induction in fluorescence intensity, while 4  
fold of induction was obtained in NF- $\kappa$ B-dlGNFP (data not shown). This  
demonstrated that the destabilized form of GNFP is applicable as  
transcription reporters.

15

**EXAMPLE 12****Construction and Functional Test for Humanized zFP506 and  
Humanized N65M**

20 Since mammalian expression is a very popular tool, human  
favored codon version is needed for better expression in mammalian  
cells. Each piece of human favored codon oligos was linked to form the  
full length of wild type and/or mutant zFP506 (hGNFP-zFP506, SEQ ID  
No. 59; hGNFP-N65M, SEQ ID No. 60). This humanized zFP506 was  
25 constituted into EGFP-N1 backbone.

Any patents or publications mentioned in this specification  
are indicative of the levels of those skilled in the art to which the  
invention pertains. These patents and publications are incorporated



reference to the same extent as if each individual publication wa  
specifically and individually indicated to be incorporated by reference.

One skilled in the art will appreciate readily that the present  
invention is adapted to carry out the objects and obtain the ends and  
5 advantages mentioned, as well as those objects and ends inherent  
therein. The present examples, along with the methods, procedures,  
treatments, molecules, and specific compounds described herein, are  
presently representative of preferred embodiments, are exemplary, and  
are not intended as limitations on the scope of the invention. Changes  
10 to the methods and compounds, and other uses, will occur to those  
skilled in the art and are encompassed within the spirit of the invention  
as defined by the scope of the claims.

**WHAT IS CLAIMED IS:**

1. A DNA sequence encoding a fluorescent protein selected from the group consisting of:

5 (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from a Class Anthozoa and wherein said organism does not exhibit bioluminescence;

(b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

10 (c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

2. The DNA sequence of claim 1, wherein said organism  
15 is from Sub-class Zoantharia.

3. The DNA sequence of claim 2, wherein said organism is from Order Zoanthidea.

20 4. The DNA sequence of claim 3, wherein said organism is from Sub-order Brachycnemina.

5. The DNA sequence of claim 4, wherein said organism is from Family Zoanthidae.

25 6. The DNA sequence of claim 5, wherein said organism is from Genus Zoanthus.

7. A DNA sequence encoding a fluorescent protein selected from the group consisting of:

(a) an isolated DNA which encodes a fluorescent protein having a nucleotide sequence selected from the group consisting of SEQ  
5 ID Nos. 55, 57-60;

(b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic  
10 code, and which encodes a fluorescent protein.

8. The DNA sequence of claim 7, wherein said DNA encodes a fluorescent protein having an amino acid sequence shown in  
SEQ ID No. 56.

15

9. The DNA sequence of claim 7, wherein said DNA is selected from the group consisting of non-humanized and humanized DNA.

20 10. The DNA sequence of claim 9, wherein said DNA is zFP506 or N65M.

11. A vector capable of expressing the DNA sequence of claim 1 in a recombinant cell, wherein said vector comprising said DNA  
25 and regulatory elements necessary for expression of the DNA in the cell.

12. The vector of claim 11, wherein said DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56.

5           13. The vector of claim 11, wherein said vector is constructed by amplifying said DNA and then inserting the amplified DNA into EGFP-N1 backbone.

10           14. The vector of claim 13, wherein said DNA is selected from the group consisting of non-humanized and humanized DNA.

15           15. The vector of claim 14, wherein said DNA is zFP506 or N65M.

          16. The vector of claim 11, wherein said vector is constructed by fusing different mouse ODC degradation domains to the C-terminal of said DNA and then inserting the fusion DNA into EGFP-N1 backbone.

20           17. The vector of claim 16, wherein said mouse ODC degradation domains are selected from the group consisting of d1, d2 and d376.

          18. The vector of claim 16, wherein said DNA is selected from the group consisting of non-humanized and humanized DNA.

25           19. The vector of claim 18, wherein said DNA is zFP506 or N65M.

20. A host cell transfected with the vector of claim 11, wherein said cell is capable of expressing a fluorescent protein.

21. The host cell of claim 20, wherein said cell is selected  
5 from the group consisting of bacterial cells, mammalian cells, plant cell, yeast and insect cells.

22. The host cell of claim 21, wherein said mammalian cell is HEK 293 cell.

10

23. The host cell of claim 21, wherein said bacterial cell is an *E. coli* cell.

24. An isolated and purified fluorescent protein coded for  
15 by DNA selected from the group consisting of:

(a) an isolated DNA which encodes a fluorescent protein from an organism from Class Anthozoa, wherein said organism does not exhibit bioluminescence;

(b) an isolated DNA which hybridizes to isolated DNA of  
20 (a) above and which encodes a fluorescent protein; and

(c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

25 25. The isolated and purified fluorescent protein of claim 24, wherein said organism is from Sub-class Zoantharia.

26. The isolated and purified fluorescent protein of claim 25, wherein said organism is from Order Zoanthidea.

27. The isolated and purified fluorescent protein of claim 26, wherein said organism is from Sub-order Brachycnemina.

5           28. The isolated and purified fluorescent protein of claim 27, wherein said organism is from Family Zoanthidae.

29. The isolated and purified fluorescent protein of claim 28, wherein said organism is from Genus Zoanthus.

10

30. An isolated and purified fluorescent protein coded for by DNA selected from the group consisting of:

(a) isolated DNA which encodes a fluorescent protein having an amino acid sequence shown in SEQ ID No. 56;

15

(b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) isolated DNA differing from said isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

20

31. The isolated and purified fluorescent protein of claim 30, wherein said protein is zFP506.

32. An amino acid sequence which can be used as a basis  
25 for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by means of hybridization, wherein said sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14.

33. The amino acid sequence of claim 32, wherein said oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16.








Figure 1






Figure 2



destabilized zfp506 data.ppt

CLOVTECH

Figure 3

1

2

3

## SEQUENCE LISTING

&lt;110&gt;

Lukyanov, Sergey A.

Labas, Yulii A.

Matz, Mikhail V.

Fradkov, Arcady F.

Jiang, Xin

Duong, Tommy

5

&lt;120&gt;

Fluorescent proteins from non-bioluminescent  
species of Class Anthozoa, genes encoding such  
proteins and uses thereof

10

&lt;130&gt;

D6196D3/PCT

&lt;141&gt;

1999-12-09

&lt;150&gt;

09/210,330

&lt;151&gt;

1998-12-11

15

&lt;160&gt;

60

&lt;210&gt;

1

&lt;211&gt;

25

&lt;212&gt;

DNA

20

&lt;213&gt;

artificial sequence

&lt;220&gt;

&lt;221&gt;

primer\_bind

&lt;223&gt;

primer TN3 used in cDNA synthesis and RACE

&lt;400&gt;

1

25 cgcagtcgac cgtttttttt ttttt

&lt;210&gt;

2

25

&lt;211&gt;

23

&lt;212&gt;

DNA

30

&lt;213&gt;

artificial sequence

&lt;220&gt;

&lt;221&gt;

primer\_bind

&lt;223&gt;

primer TS used in cDNA synthesis and RACE

&lt;400&gt;

2

35 aagcagtggt atcaacgcag agt

&lt;210&gt;

3

23

&lt;211&gt;

6

&lt;212&gt;

PRT





5      <213>      *Aequorea victoria*  
      <220>  
      <222>      21  
      <223>      amino acid sequence of a key stretch on which  
                  primer NGH is based; Xaa at position 21  
                  represents unknown

<400>      3

Gly Xaa Val Asn Gly His  
5

10

<210>      4

<211>      20

<212>      DNA

<213>      artificial sequence

<220>

<221>      primer\_bind

<222>      12

<223>      primer NGH used for isolation of fluorescent  
protein; n at position 12 represents any of the  
four bases

<400>      4

gayggctgcg tnaayggdca

20

<210>      5

<211>      5

<212>      PRT

<213>      *Aequorea victoria*

<220>

<222>      31...35

<223>      amino acid sequence of a key stretch on which  
primers GEGa and GEGb are based


<400>      5

Gly Glu Gly Glu Gly  
5

35

<210>      6

<211>      20



<212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 5 <223> primer GEGa used for isolation of fluorescent  
 protein  
 <400> 6

gttacaggtg arggmgargg

20

10 <210> 7  
 <211> 20  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 15 <221> primer\_bind  
 <223> primer GEGb used for isolation of fluorescent  
 protein  
 <400> 7

gttacaggtg arggkgargg

20

20 <210> 8  
 <211> 5  
 <212> PRT  
 <213> *Aequorea victoria*  
 25 <220>  
 <222> 31...35  
 <223> amino acid sequence of a key stretch on which  
 primers GNGa and GNGb are based  
 <400> 8

30 Gly Glu Gly Asn Gly  
 5

<210> 9  
 <211> 20  
 35 <212> DNA  
 <213> artificial sequence  
 <220>



<221> primer\_bind  
 <223> primer GNGa used for isolation of fluorescent  
 protein  
 <400> 9

5 gttacaggtg arggmaaygg 20

<210> 10  
 <211> 20  
 <212> DNA  
 10 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> primer GNGb used for isolation of fluorescent  
 protein  
 15 <400> 10

gttacaggtg arggkaaygg 20

<210> 11  
 <211> 5  
 20 <212> PRT  
 <213> *Aequorea victoria*  
 <220>  
 <222> 127...131  
 <223> amino acid sequence of a key stretch on which  
 25 primer NFP is based  
 <400> 11

Gly Met Asn Phe Pro  
 5

30 <210> 12  
 <211> 5  
 <212> PRT  
 <213> *Aequorea victoria*  
 <220>  
 35 <222> 127...131  
 <223> amino acid sequence of a key stretch on which  
 primer NFP is based



&lt;400&gt; 12

Gly Val Asn Phe Pro

5

5 <210> 13  
 <211> 20  
 <212> DNA  
 <213> artificial sequence  
 <220>

10 <221> primer\_bind  
 <223> primer NFP used for isolation of fluorescent  
 protein

&lt;400&gt; 13

15 ttccayggtr tgaayttycc

20

<210> 14  
 <211> 4  
 <212> PRT  
 <213> *Aequorea victoria*  
 <220>

20 <222> 134...137  
 <223> amino acid sequence of a key stretch on which  
 primers PVMa and PVMb are based

&lt;400&gt; 14

25 Gly Pro Val Met

<210> 15  
 <211> 21  
 <212> DNA  
 <213> artificial sequence  
 <220>

30 <221> primer\_bind  
 <222> 15

35 <223> primer PVMa used for isolation of fluorescent  
 protein; n at position 15 represents any of the





four bases  
 <400> 15  
 cctgccrayg gtcnngtmat g 21

5 <210> 16  
 <211> 21  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 10 <221> primer\_bind  
 <222> 15  
 <223> primer PVMb used for isolation of fluorescent  
 protein; n at position 15 represents any of the  
 four bases  
 15 <400> 16  
 cctgccrayg gtcnngtkat g 21

<210> 17  
 <211> 47  
 20 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> primer T7-TN3 used in cDNA synthesis and RACE  
 25 <400> 17  
 gtaatacgac tcactatagg gccgcagtcg accgtttttt tttttt 47

<210> 18  
 <211> 45  
 30 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> primer T7-TS used in cDNA synthesis and RACE  
 35 <400> 18



gtaatacgac tcactatagg gcaagcagtg gtatcaacgc agagt 45  
    <210> 19  
    <211> 22  
    <212> DNA  
5     <213> artificial sequence  
    <220>  
    <221> primer\_bind  
    <223> primer T7 used in cDNA synthesis and RACE  
    <400> 19

10 gtaatacgac tcactatagg gc 22  
    <210> 20  
    <211> 21  
    <212> DNA  
15     <213> artificial sequence  
    <220>  
    <221> primer\_bind  
    <223> gene-specific primer used for 5'-RACE for  
          *Anemonia majano*  
20     <400> 20

gaaatagtca ggcatactgg t 21  
    <210> 21  
    <211> 20  
25     <212> DNA  
    <213> artificial sequence  
    <220>  
    <221> primer\_bind  
    <223> gene-specific primer used for 5'-RACE for  
          *Anemonia majano*  
30     <400> 21

gtcaggcata ctggtaggat 20  
    <210> 22  
35     <211> 21

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200

201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300

<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
5 <223> gene-specific primer used for 5'-RACE for  
Clavularia sp.  
<400> 22  
cttgaaatag tctgctatat c 21

10 <210> 23  
<211> 19  
<212> - DNA  
<213> artificial sequence  
<220>  
15 <221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
Clavularia sp.  
<400> 23  
tctgctatat cgtctgggt 19

20  
<210> 24  
<211> 23  
<212> DNA  
<213> artificial sequence  
25 <220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
Zoanthus sp.  
<400> 24

30 gttcttgaaa tagtctacta tgt 23  
  
<210> 25  
<211> 20  
<212> DNA  
35 <213> artificial sequence

com.

2023-09-01

10/10/2023

SEQ 9/23





*Discosoma striata*

&lt;400&gt; 28

ttgtcttctt ctgcacaac 19

5 &lt;210&gt; 29

&lt;211&gt; 17

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

10 &lt;221&gt; primer\_bind

<223> gene-specific primer used for 5'-RACE for  
*Discosoma striata*

&lt;400&gt; 29

ctgcacaacg ggtccat 17

15

&lt;210&gt; 30

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

20 &lt;220&gt;

&lt;221&gt; primer\_bind

<223> gene-specific primer used for 5'-RACE for  
*Anemonia sulcata*

&lt;400&gt; 30

25 cctctatctt catttcctgc 20

&lt;210&gt; 31

&lt;211&gt; 20

&lt;212&gt; DNA

30 &lt;213&gt; artificial sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

<223> gene-specific primer used for 5'-RACE for  
*Anemonia sulcata*

35 &lt;400&gt; 31

tatcttcatt tcctgcgtac 20



	<210>	32	
	<211>	19	
	<212>	DNA	
5	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
	<223>	gene-specific primer used for 5'-RACE for	
		<i>Discosoma</i> sp. "magenta"	
10	<400>	32	
		ttcagcaccc catcacgag	19
	<210>	33	
	<211>	19	
15	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
	<223>	gene-specific primer used for 5'-RACE for	
20		<i>Discosoma</i> sp. "magenta"	
	<400>	33	
		acgctcagag ctgggttcc	19
	<210>	34	
25	<211>	22	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
30	<223>	gene-specific primer used for 5'-RACE for	
		<i>Discosoma</i> sp. "green"	
	<400>	34	
		ccctcagcaa tccatcacgt tc	22
35	<210>	35	
	<211>	20	
	<212>	DNA	



<213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> gene-specific primer used for 5'-RACE for  
 5 *Discosoma* sp. "green"  
 <400> 35  
 attatctcag tggatgggtc 20  
 <210> 36  
 10 <211> 31  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 15 <223> upstream primer used to obtain full coding region  
 of nFPs from *Anemonia majano*  
 <400> 36  
 acatggatcc gctctttcaa acaagtttat c 31  
 20 <210> 37  
 <211> 34  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 25 <221> primer\_bind  
 <223> downstream primer used to obtain full coding  
 region of nFPs from *Anemonia majano*  
 <400> 37  
 tagtactcga gcttattcgt atttcagtga aatc 34  
 30 <210> 38  
 <211> 29  
 <212> DNA  
 <213> artificial sequence  
 35 <220>  
 <221> primer\_bind  
 <223> upstream primer used to obtain full coding region



of nFPs from *Clavularia sp.*

<400> 38

acatggatcc aacatttttt tgagaaacg 29

5 <210> 39

<211> 28

<212> DNA

<213> artificial sequence

<220>

10 <221> primer\_bind

<223> upstream primer used to obtain full coding region  
of nFPs from *Clavularia sp.*

<400> 39

acatggatcc aaagctctaa ccaccatg 28

15 <210> 40

<211> 31

<212> DNA

<213> artificial sequence

20 <220>

<221> primer\_bind

<223> downstream primer used to obtain full coding  
region of nFPs from *Clavularia sp.*

<400> 40

25 tagtactcga gcaacacaaa ccctcagaca a 31

<210> 41

<211> 28

<212> DNA

30 <213> artificial sequence

<220>

<221> primer\_bind

<223> upstream primer used to obtain full coding region  
of nFPs from *Zoanthus sp.*

35 <400> 41

acatggatcc gctcagtcaa agcacggt 28





<210> 42  
 <211> 32  
 <212> DNA  
 5 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> downstream primer used to obtain full coding  
 region of nFPs from *Zoanthus sp.*  
 10 <400> 42  
 tagtactcga ggttggaact acattcttat ca 32  
  
 <210> 43  
 <211> 31  
 15 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> upstream primer used to obtain full coding region  
 20 of nFPs from *Discosoma sp.* "red"  
 <400> 43  
 acatgatcc aggtcttcca agaatgttat c 31  
  
 <210> 44  
 25 <211> 29  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 30 <223> downstream primer used to obtain full coding  
 region of nFPs from *Discosoma sp.* "red"  
 <400> 44  
 tagtactcga ggagccaagt tcagcctta 29  
  
 35 <210> 45  
 <211> 28  
 <212> DNA



<213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> upstream primer used to obtain full coding region  
 5 of nFPs from *Discosoma striata*  
 <400> 45  
 acatggatcc agttggtcca agagtgtg 28  
 <210> 46  
 10 <211> 28  
 <212> DNA  
 <213> \_ artificial sequence  
 <220>  
 <221> primer\_bind  
 15 <223> downstream primer used to obtain full coding  
 region of nFPs from *Discosoma striata*  
 <400> 46  
 tagcgagctc tatcatgcct cgtcacct 28  
 20 <210> 47  
 <211> 31  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 25 <221> primer\_bind  
 <223> upstream primer used to obtain full coding region  
 of nFPs from *Anemonia sulcata*  
 <400> 47  
 acatggatcc gcttcctttt taaagaagac t 31  
 30 <210> 48  
 <211> 28  
 <212> DNA  
 <213> artificial sequence  
 35 <220>  
 <221> primer\_bind  
 <223> downstream primer used to obtain full coding



region of nFPs from *Anemonia sulcata*

<400> 48

tagtactcga gtccttggga gcggttg 28

5 <210> 49

<211> 30

<212> DNA

<213> artificial sequence

<220>

10 <221> primer\_bind

<223> upstream primer used to obtain full coding region  
of nFPs from *Discosoma sp. "magenta"*

<400> 49

acatggatcc agttgttcca agaatgtgat 30

15

<210> 50

<211> 26

<212> DNA

<213> artificial sequence

20 <220>

<221> primer\_bind

<223> downstream primer used to obtain full coding  
region of nFPs from *Discosoma sp. "magenta"*

<400> 50

25 tagtactcga ggccattacg ctaatc 26

<210> 51

<211> 31

<212> DNA

30 <213> artificial sequence

<220>

<221> primer\_bind

<223> upstream primer used to obtain full coding region  
of nFPs from *Discosoma sp. "green"*

35 <400> 51

acatggatcc agtgcactta aagaagaaat g 31



<210> 52  
 <211> 29  
 <212> DNA  
 5 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> downstream primer used to obtain full coding  
 region of nFPs from *Discosoma* sp. "green"  
 10 <400> 52

tagtactcga gattcgggttt aatgccttg 29

<210> 53  
 <211> 33  
 15 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> TS-oligo used in cDNA synthesis and RACE  
 20 <400> 53

aagcagtgggt atcaacgcag agtacgcrgr grg 33

<210> 54  
 <211> 238  
 25 <212> PRT  
 <213> *Aequorea victoria*  
 <220>  
 <223> amino acid sequence of GFP  
 <400> 54

30 Met Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu  
                                   5                                  10                                  15  
 Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser  
                                   20                                  25                                  30  
 Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys  
 35                                  35                                  40                                  45  
 Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu  
                                   50                                  55                                  60





	Val	Thr	Thr	Phe	Ser	Tyr	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro
					65					70					75
	Asp	His	Met	Lys	Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro	Glu
					80					85					90
5	Gly	Tyr	Val	Gln	Glu	Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn
					95					100					105
	Tyr	Lys	Thr	Arg	Ala	Glu	Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val
					110					115					120
	Asn	Arg	Ile	Glu	Leu	Lys	Gly	Ile	Asp	Phe	Lys	Glu	Asp	Gly	Asn
10					125					130					135
	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr	Asn	Tyr	Asn	Ser	His	Asn	Val
					140					145					150
	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	Gly	Ile	Lys	Val	Asn	Phe
					155					160					165
15	Lys	Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	Val	Gln	Leu	Ala	Asp
					170					175					180
	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly	Pro	Val	Leu	Leu
					185					190					195
	Pro	Asp	Asn	His	Tyr	Leu	Ser	Thr	Gln	Ser	Ala	Leu	Ser	Lys	Asp
20					200					205					210
	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu	Leu	Glu	Phe	Val	Thr
					215					220					225
	Ala	Ala	Gly	Ile	Thr	His	Gly	Met	Asp	Glu	Leu	Tyr	Lys		
					230					235					

25

&lt;210&gt; 55

&lt;211&gt; 858

&lt;212&gt; DNA

<213> *Zoanthus* sp.

30

&lt;220&gt;

&lt;221&gt; CDS

&lt;223&gt; cDNA sequence of zFP506

&lt;400&gt; 55

cccaaagcta ctcgactttt gtatcacctt tggggcatca agcgatctgt 50  
 35 tctcaacatg gctcagtcaa agcacggtct aacaaaagaa atgacaatga 100  
 aataccgtat ggaaggggtgc gtcgatggac ataaatttgt gatcacggga 150  
 gagggcattg gatatccgtt caaagggaaa caggctatta atctgtgtgt 200  
 ggtcgaaggt ggaccattgc catttgccga agacatattg tcagctgcct 250



```

    ttaactacgg aaacaggggtt ttactgaat atcctcaaga catagttgac 300
    tatttcaaga actcgtgtcc tgctggatat acatgggaca ggtcttttct 350
    ctttgaggat ggagcagttt gcatatgtaa tgcagatata acagtgagtg 400
    ttgaagaaaa ctgcatgtat catgagtcca aattttatgg agtgaatttt 450
5   cctgctgatg gacctgtgat gaaaaagatg acagataact gggagccatc 500
    ctgcgagaag atcataccag tacctaagca ggggatattg aaaggggatg 550
    tctccatgta cctccttctg aaggatgggtg ggcgtttacg gtgccaattc 600
    gacacagttt acaaagcaaa gtctgtgcca agaaagatgc cggactggca 650
    cttcatccag cataagctca cccgtgaaga ccgcagcgat gctaagaatc 700
10  agaaatggca tctgacagaa catgctattg catccggatc tgcattgccc 750
    tgataagaat gtagctccag cattctaattg catgtgcttg ccaattattc 800
    tgataaatat gtagttgagt tggaaacaga ctagtacaaa taaaagcaca 850
    tttaaatac                                     858

```

```

15      <210>      56
          <211>      230
          <212>      PRT
          <213>      Zoanthus sp.
          <220>
20      <223>      amino acid sequence of zFP506
          <400>      56
Ala Gln Ser Lys His Gly Leu Thr Lys Glu Met Thr Met Lys Tyr
          5                                10                                15
Arg Met Glu Gly Cys Val Asp Gly His Lys Phe Val Ile Thr Gly
25          20                                25                                30
Glu Gly Ile Gly Tyr Pro Phe Lys Gly Lys Gln Ala Ile Asn Leu
          35                                40                                45
Cys Val Val Glu Gly Gly Pro Leu Pro Phe Ala Glu Asp Ile Leu
          50                                55                                60
30  Ser Ala Ala Phe Asn Tyr Gly Asn Arg Val Phe Thr Glu Tyr Pro
          65                                70                                75
Gln Asp Ile Val Asp Tyr Phe Lys Asn Ser Cys Pro Ala Gly Tyr
          80                                85                                90
Thr Trp Asp Arg Ser Phe Leu Phe Glu Asp Gly Ala Val Cys Ile
35          95                                100                               105
Cys Asn Ala Asp Ile Thr Val Ser Val Glu Glu Asn Cys Met Tyr
          110                               115                               120

```



[illegible]

	<210>	57
	<211>	826
20	<212>	DNA
	<213>	artificial sequence
	<220>	
	<222>	497
	<223>	nucleic acid sequence of d1-zFP506, n at position
25		497 represents unknown
	<400>	57

	atggctcagt	caaagcacgg	tctaacaaaa	gaaatgacaa	tgaaataccg
	tatggaaggg	tgcgtcgatg	gacataaatt	tgtgatcacg	ggagagggca
	ttggatatcc	gttcaaaggg	aaacaggcta	ttaatctgtg	tgtgggtcgaa
30	ggtggaccat	tgccatttgc	cgaagacata	ttgtcagctg	cctttaacta
	cggaaacagg	gttttctactg	aatatcctca	agacatagtt	gactatttca
	agaactcgtg	tcttgcctgga	tatacatggg	acagggtcttt	tctctttgag
	gatggagcag	tttgcatatg	taatgcagat	ataacagtga	gtgttggaaga
	aaactgcgatg	tatcatgagt	ccaaattttta	tggagtgaat	tttctctgctg
35	atggacctgt	gatgaaaaag	atgacagata	actgggagcc	atcctgcgag
	aagatcatac	cagtacctaa	gcaggggata	ttgaaagggg	atgtctncat
	gtacctcctt	ctgaagggatg	gtgggctgttt	acggtgccaa	ttcgacacag



tttacaaagc aaagtctgtg ccaagaaaga tgccggactg gcacttcac  
 cagcataagc tcacccgtga agaccgcagc gatgctaaga atcagaaatg  
 gcatctgaca gaacatgcta ttgcatccgg atctgcattg cccaagctta  
 gccatggctt cccgccggcg gtggcgggcg aggatgatgg cacgctgccc  
 5 atgtcttgtg cccaggagag cgggatggac cgtcaccctg cagcctgtgc  
 ttctgctagg atcaatgtgt agatgc 826

<210> 58  
 <211> 826  
 10 <212> DNA  
 <213> artificial sequence  
 <220> —  
 <222> 497  
 <223> nucleic acid sequence of d2-zFP506, n at position  
 15 497 represents unknown

<400> 58  
 atggctcagt caaagcacgg tctaacaaaa gaaatgacaa tgaaataccg 50  
 tatggaaggg tgcgtcgatg gacataaatt tgtgatcacg ggagaggggca 100  
 ttggatatcc gttcaaaggg aaacaggcta ttaatctgtg tgtggtcgaa 150  
 20 ggtggaccat tgccatttgc cgaagacata ttgtcagctg cctttaacta 200  
 cggaaacagg gttttcactg aatatcctca agacatagtt gactatttca 250  
 agaactcgtg tcctgctgga tatacatggg acaggctctt tctctttgag 300  
 gatggagcag tttgcatatg taatgcagat ataacagtga gtggtgaaga 350  
 aaactgcatg tatcatgagt ccaaatttta tggagtgaat tttcctgctg 400  
 25 atggacctgt gatgaaaaag atgacagata actgggagcc atcctgagag 450  
 aagatcatat cagtacctaa gcaggggata ttgaaagggg atgtctncat 500  
 gtacctcctt ctgaaggatg gtgggctgtt acggtgccaa ttcgacacag 550  
 tttacaaagc aaagtctgtg ccaagaaaga tgccggactg gcacttcac 600  
 cagcataagc tcacccgtga agaccgcagc gatgctaaga atcagaaatg 650  
 30 gcatctgaca gaacatgcta ttgcatccgg atctgcattg cccaagctta 700  
 gccatggctt cccgccggag gtggaggagc aggatgatgg cacgctgccc 750  
 atgtcttgtg cccaggagag cgggatggac cgtcaccctg cagcctgtgc 800  
 ttctgctagg atcaatgtgt agatgc 826

35 <210> 59  
 <211> 696  
 <212> DNA





<213> artificial sequence

<220>

<223> nucleic acid sequence of humanized zFP506

<400> 59

```

5  atggcccagt ccaagcacgg cctgaccaag gagatgacca tgaagtaccg  50
   catggagggc tgcgtggacg gccacaagtt cgtgatcacc ggcgagggca 100
   tcggctaccc cttcaagggc aagcaggcca tcaacctgtg cgtggtggag 150
   ggcggcccct tgccttcgc cgaggacatc ttgtccgccg ccttcaacta 200
   cggcaaccgc gtgttcaccg agtaccacca ggacatcgtc gactacttca 250
10 agaactcctg ccccgccggc tacacctggg accgctcctt cctgttcgag 300
   gacggcgccg tgtgcatctg caacgccgac atcacctga gcgtggagga 350
   gaactgcatg taccacgagt ccaagttcta cggcgtgaac ttccccgccg 400
   acggccccgt gatgaagaag atgaccgaca actgggagcc ctctgcgag 450
   aagatcatcc ccgtgccc aa gcagggcac ttgaaggcg acgtgagcat 500
15 gtacctgctg ctgaaggacg gtggccgctt gcgctgccag ttcgacaccg 550
   tgtacaaggc caagtccgtg ccccgcaaga tgcccgactg gcacttcac 600
   cagcacaagc tgaccgcga ggaccgcagc gacgccaaga accagaagtg 650
   gcacctgacc gagcacgcca tcgcctccgg ctccgccttg ccctga      696

```

20 <210> 60

<211> 696

<212> DNA

<213> artificial sequence

<220>

25 <223> nucleic acid sequence of humanized N65M

<400> 60

```

   atggcccagt ccaagcacgg cctgaccaag gagatgacca tgaagtaccg  50
   catggagggc tgcgtggacg gccacaagtt cgtgatcacc ggcgagggca 100
   tcggctaccc cttcaagggc aagcaggcca tcaacctgtg cgtggtggag 150
30 ggcggcccct tgccttcgc cgaggacatc ttgtccgccg ccttcatgta 200
   cggcaaccgc gtgttcaccg agtaccacca ggacatcgtc gactacttca 250
   agaactcctg ccccgccggc tacacctggg accgctcctt cctgttcgag 300
   gacggcgccg tgtgcatctg caacgccgac atcacctga gcgtggagga 350
   gaactgcatg taccacgagt ccaagttcta cggcgtgaac ttccccgccg 400
35 acggccccgt gatgaagaag atgaccgaca actgggagcc ctctgcgag 450
   aagatcatcc ccgtgccc aa gcagggcac ttgaaggcg acgtgagcat 500
   gtacctgctg ctgaaggacg gcggccgctt gcgctgccag ttcgacaccg 550

```



tgtacaaggc caagtccgtg ccccgcaaga tgcccgactg gcacttcac 600  
cagcacaagc tgaccgcga ggaccgcagc gacgccaaga accagaagtg 650  
gcacctgacc gagcacgcca tcgcctccgg ctccgccttg ccctga 696



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29291

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) Please See Extra Sheet.

US CL Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/320.1, 252.3, 252.33, 325, 410, 254.11, 348, 369, 69.1; 530/350; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	MATZ et al. Fluorescent proteins from nonbioluminescent Anthozoa species. Nature Biotechnology. October 1999, Volume 17, No. 10, pages 969-673, entire document.	1-33
X, P	DE 197 18 640 A1 (WIEDENMANN) 22 July 1999, entire document.	24-25, 30

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* Later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* Document defining the general state of the art which is not considered to be of particular relevance	*X* Document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* Earlier document published on or after the international filing date	*Y* Document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* Document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* Document member of the same patent family
*O* Document referring to an oral disclosure, use, exhibition or other means	
*P* Document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

23 FEBRUARY 2000

Date of mailing of the international search report

17 MAR 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GABRIELE ELISABETH BUGAISKY

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29291

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANDERLUH et al. Cloning, sequencing, and expression of equinatoxin II. Biochemical and Biophysical Research Communications. 1996, Volume 220, No. 2, pages 437-442, entire document.	1-2, 7, 11, 20-21, 23-25, 30
X — A —	MACEK et al. Intrinsic tryptophan fluorescence of equinatoxin II, a pore-forming polypeptide from the sea anemone, Actinia equina L, monitors its interaction with lipid membranes. European Journal of Biochemistry. 1995, Volume 234, pages 329-335, entire document.	24-25, 30 — 1-2, 7, 11, 20-21, 23

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/29291

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

C07K 14/435; C12N 1/00, 1/15, 1/21, 5/10, 15/12, 15/63

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/ 320.1, 252.3, 252.33, 325, 410, 254.11, 348, 369; 530/350; 536/23.5

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

Dialog files 155, 5, 434, 34, 44, 357, 35 (Medline, Biosis, Scisearch, Oceanic Abs., Derwent Biotech Abs, Dissertation Abs.), STN-CAS files Registry, CAPLUS; WEST, files USPT, Derwent WPI  
search terms: fluoresc?, bioluminesc?, protein, anthozo?, Zoanth?, Corralimorph?, Discosom?, Coral?, alga, algae, Cnidar?, Invert?, Rhodact?, Actinodisc?, magenta, Clavularia, Zoanthus, Anemonia, majano, anemon?, Zoanthar?, Actinar?, Zoanthid?, Stolonif?, Alcyonar? aqskhgktke/sqsp, zfp506, striata, sulcata, brachycnem?, Isaurus

